

## COMMENTARY

# Bridging the Gap Between Spatial and Mnemonic Views of the Hippocampal Formation

Elizabeth A. Buffalo\*

**ABSTRACT:** While it has long been recognized that medial temporal lobe structures are important for memory formation, studies in rodents have also identified exquisite spatial representations in these regions in the form of place cells in the hippocampus and grid cells in the entorhinal cortex. Spatial representations entail neural activity that is observed when the rat is in a given physical location, and these representations are thought to form the basis of navigation via path integration. Recent studies in nonhuman primates have suggested that similar kinds of spatial representations can be identified, even in the absence of physical movement through an environment. Here, I will highlight some recent work that addresses similarities and differences between spatial responses as identified in rodents and primates. I will also discuss areas of opportunity for future research to further our understanding of the function of the hippocampal formation. © 2015 Wiley Periodicals, Inc.

**KEY WORDS:** monkey; memory; hippocampus; entorhinal cortex; navigation

## INTRODUCTION

The ability of the mammalian brain to store and later retrieve information is remarkable. Detailed, complex memories can be formed after as little as one exposure, and those memories can be retained for decades. This ability is compromised following damage to structures located in the medial temporal lobe, including the hippocampus and surrounding cortical regions (Squire and Zola-Morgan, 1991). While it has long been recognized that these structures are important for memory (Eichenbaum et al., 1996; Suzuki et al., 1997; Suzuki and Eichenbaum, 2000; Wirth et al., 2003; Rutishauser et al., 2006, 2010; Jutras et al., 2009; Yanike et al., 2009; Jutras and Buffalo, 2010; Naya and Suzuki, 2011), a largely parallel line of research in rodents has highlighted the contribu-

tion of these same structures to our sense of space (O'Keefe and Dostrovsky, 1971; O'Keefe, 1976; O'Keefe and Nadel, 1978; Rolls et al., 1989; Ekstrom et al., 2003; Hafting et al., 2005; Sargolini et al., 2006; Fyhn et al., 2008; Moser et al., 2008; Doeller et al., 2010). While these two perspectives on the function of the hippocampal formation, e.g., important for memory or providing an internal map, have fueled research for decades, we are still in the early stages of reconciling these two views. In that endeavor, studies with the nonhuman primate provide an important opportunity to bridge the gap between neurophysiological studies of spatial coding carried out largely in rodents and behavioral studies in human amnesic patients. In this commentary, I will discuss some recent findings from nonhuman primates that were inspired by the findings of robust spatial coding in the rodent hippocampal formation, and I will describe future areas of opportunity to advance our understanding of the hippocampal formation.

## ARE SPATIAL REPRESENTATIONS IN THE HIPPOCAMPAL FORMATION SIMILAR ACROSS SPECIES?

The existence of spatial representations in the hippocampal formation has been appreciated since the truly groundbreaking work of John O'Keefe in the early 1970s. O'Keefe et al. demonstrated the existence of place cells in the rodent hippocampus (O'Keefe and Dostrovsky, 1971; O'Keefe, 1976; O'Keefe and Nadel, 1978). Place cells are neurons that fire action potentials whenever the rat is in a specific place in an environment, the neuron's place field. The combined activity of many of these neurons, with distinct place fields, effectively provide a map of the environment and, in more recent research, it was demonstrated that the rat's trajectory through space can be accurately decoded by measuring the activity of these neurons (Jensen and Lisman, 2000). Place cells with the sharpest and most reliable place fields are found in the dorsal part of the rodent hippocampus (O'Keefe and Nadel, 1978; McNaughton et al., 2006). To

Department of Physiology and Biophysics, University of Washington, Seattle, Washington

Grant sponsor: National Institutes of Health; Grant numbers: MH080007; MH 093807; Grant sponsor: Office of Infrastructure Programs (ORIP); Grant number: P51OD010425.

\*Correspondence to: Elizabeth A. Buffalo, Department of Physiology and Biophysics, University of Washington, Seattle, WA, USA.

E-mail: ebuffalo@uw.edu

Accepted for publication 17 March 2015.

DOI 10.1002/hipo.22444

Published online 18 March 2015 in Wiley Online Library (wileyonlinelibrary.com).

understand what gives rise to these spatial representations, May-Britt and Edvard Moser began recording in the dorsolateral band of the medial entorhinal cortex, the part of the rodent brain that provides the strongest input to the dorsal hippocampus. Through this work, they identified periodic spatial representations that they called entorhinal grid cells (Fyhn et al., 2004; Hafting et al., 2005). Like place cells, grid cells represent the location of the rat, but each grid cell has multiple place fields. The amazing thing about grid cells is that the multiple place fields lie in precise geometric relation to each other and form a tessellated array of equilateral triangles, a “grid” that tiles the environment. Accordingly, a spatial autocorrelation of the grid field map produces a hexagonal structure, with 60° rotational symmetry.

While there is a large body of literature describing spatial representations in the hippocampal formation in rodents (Moser et al., 2008), relatively little is known about similar representations in the nonhuman primate (Rolls et al., 1997). One striking difference between rodents and primates is the way in which information about the external world is obtained. Rodents typically gather information by moving to visit different locations in the environment, sniffing and whisking. By contrast, primates chiefly use eye movements to visually explore an environment, and our visual system allows for inspection of the environment at a distance. Accordingly, a rodent’s exploration through space may use the same neural coding mechanisms as a primate’s visual exploration. Indeed, while grid cells in rodents and bats (Yartsev et al., 2011) have been identified as animals move about and explore an environment, we recently identified entorhinal grid cells in primates that represent space during visual exploration (Killian et al., 2012). As monkeys freely viewed complex natural images on a computer monitor, these cells responded whenever the monkey fixated specific regions of the monitor, independent of the stimulus content. The locations of fixation that evoked activity in an individual neuron formed the nodes of equilateral triangles that tiled the computer monitor, with hexagonal tiling in the autocorrelation. This study provided the first direct evidence for the existence of grid cells in the primate, and, importantly, suggested that grid cells may encode spatial locations even in a non-moving animal. These data are consistent with early work demonstrating hippocampal “view cells,” cells that responded whenever a monkey looked toward specific locations in the environment (Rolls, 1999). Subsequent studies investigated spatial representations in the human hippocampal formation by recording from epilepsy patients with surgically implanted electrodes as the patients navigated through virtual environments. These studies revealed place cells in the human hippocampus (Miller et al., 2013) as well as grid cells in the human entorhinal cortex (Jacobs et al., 2013).

Other key components of the rodent hippocampal navigation system include head direction cells, cells with a heading preference when the rodent is within a particular environment (Taube et al., 1990; Taube, 1995; Sargolini et al., 2006; Boccaro et al., 2010), and border or boundary vector cells, cells that are selectively active at specific distances from environmen-

tal boundaries (Solstad et al., 2008; Lever et al., 2009). In the primate, putative border cells have been identified, cells that are active whenever the monkey fixates one of the edges of the visual image (Killian et al., 2012), along with “saccade direction” cells, cells that demonstrate selectivity for the angle of the upcoming or just-completed saccade (Killian et al., 2013). Given the potential analogy between visual exploration in primates and spatial exploration in rodents, saccade direction cells may provide directional information similar to head direction cells. The direction of saccades carries important information about past and future gaze locations. Notably, recent research in humans using virtual navigation has identified novel spatial representations, not previously identified in the rodent. In particular, neurons in the human entorhinal cortex have been identified that encode the direction of heading in a circular environment, i.e., clockwise or counter-clockwise (Jacobs et al., 2010). Taken together, these findings provide evidence that a full range of spatial representations exist in the primate hippocampal formation. Through experiments with visual exploration and virtual navigation, these studies present a new research avenue for understanding and comparing neuronal representations across species. However, they also raise important questions about the coordinate frame for these spatial representations, the extent to which these representations reflect visuospatial versus environmental navigation, and the interaction between spatial and mnemonic representations. These questions drive current and future research in this area.

### DOES THE HIPPOCAMPAL FORMATION PERFORM SIMILAR COMPUTATIONS ACROSS SPECIES?

The perspective that emphasizes the importance of medial temporal lobe structures to memory began in the 1950s with the report of the devastating memory impairment observed in patient H.M. following medial temporal resection to relieve otherwise intractable epilepsy (Scoville and Milner, 1957). This tradition continued through decades of research with other amnesic patients (Squire and Zola-Morgan, 2011) along with work on the nonhuman primate model of amnesia (Mishkin, 1982; Zola-Morgan et al., 1982). A fundamental understanding derived from these studies is that damage to medial temporal lobe structures produces impairment in long-term memory, but leaves short-term or working memory intact. Working memory refers to memory that is short-lasting and typically involves rehearsal; the content of working memory is information that is available to conscious awareness as long as the subject continues to rehearse it. Working memory is thought to be entirely independent of medial temporal lobe structures (Drachman and Arbib, 1966; Baddeley and Warrington, 1970; Milner, 1972; Squire et al., 2004; Jenson et al., 2012). In this regard, path integration provides an interesting point of comparison between the spatial and memory perspectives of the

hippocampal formation. Path integration refers to the ability to keep track of one's position in space and the ability to navigate towards a reference location by way of self-motion cues. One strong hypothesis about the function of spatial representations in the hippocampal formation is that the path integrator is located *within* the entorhinal cortex (McNaughton et al., 2006). If that is true, then one would predict that damage to the entorhinal cortex would produce an impairment in path integration. However, path integration refers to an on-line computation of spatial position, and as such, falls into the realm of working memory. This hypothesis is therefore inconsistent with the distinction between short term and long term memory in terms of reliance upon medial temporal lobe structures. According to the memory tradition, in the presence of damage to the hippocampal formation, short-term memory is intact, and performance should be intact as long as information is able to be held in mind, even if that information is spatial. According to the spatial tradition, there might be an exception to this idea because the entorhinal cortex is needed for on-line computations involved in path integration.

In rodents, several studies have demonstrated impairment in path integration following lesions of the hippocampus or entorhinal cortex (Maaswinkel et al., 1999; Parron and Save, 2004; Kim et al., 2013), but see (Alyan and McNaughton, 1999). By contrast, amnesic patients with bilateral damage restricted to the hippocampus or involving the hippocampus plus adjacent cortical structures showed completely intact path integration relative to control subjects (Shrager et al., 2008; Kim et al., 2013). Notably, patients with medial temporal lobe damage performed worse when demands on long-term memory were increased by introducing distraction (Shrager et al., 2008). Together, these data suggest that in humans, the hippocampus and entorhinal cortex are not unique sites that are essential for path integration because patients perform normally as long as the route is fairly simple and can be held in working memory. It is possible that the computations needed for spatial navigation are accomplished in regions upstream from the medial temporal lobe, and that these computations arrive from cortex just as information from other modalities reaches the medial temporal lobe (Suzuki and Amaral, 1994). By this view, the critical role of the medial temporal lobe is in carrying out the operation of transforming online perceptual information into long-term memory. By contrast, rodents with damage to the hippocampal formation are impaired, even on very short paths, i.e., traveling less than 1 m and within 3 turns (Kim et al., 2013), suggesting that spatial computations that support path integration are carried out online in the hippocampal formation. There are several ways to understand these inconsistencies across species. It is possible that rodents and humans use different strategies when solving these tasks, with humans relying more on visual imagery rather than self-motion cues. It is also possible that the capacity of working memory is different for humans and rodents (Kim et al., 2013). Finally, it is possible that the hippocampal formation has evolved to perform different computations in humans from rodents. While it is currently technically challenging to perform studies of path

integration in freely-moving monkeys, it is possible that future experiments combining targeted lesions in monkeys with path integration performed in virtual reality (Tcheang et al., 2011; Sherrill et al., 2013) could advance our understanding of similarity of function between the rodent and primate hippocampal formation.

## FUTURE AREAS OF OPPORTUNITY

The demonstration of exquisite spatial representations in the form of place cells and grid cells have spawned decades of research that has advanced our understanding of the development of these representations (Tan et al., 2015; Langston et al., 2010; Wills et al., 2010) and potential mechanisms for their generation (McNaughton et al., 2006; Burgess, 2008; Welinder et al., 2008). However, what is clearly a significant gap in our understanding concerns their function. The original idea, as put forward by O'Keefe and Nadel in 1978, is that these kinds of spatial representations serve as a "cognitive map," an internal metric for our location, derived from the integration of environmental sensory information with self-motion information. As discussed above, the idea of a "cognitive map" leans heavily on the notion of performing spatial computations as the critical function of the hippocampus, and the idea of the entorhinal cortex as an odometer has followed from this perspective (Jeffery, 2013). In contrast, even the first report of grid cells raised the suggestion that spatial representations may provide an important substrate for memory formation:

"... the well-established role of the hippocampus in spatial navigation may reflect the essential nature of spatial input as an element of most episodic memories rather than a specific role in computing the animal's location within a given context. Rather than calculating location per se, hippocampal networks may transform spatial and nonspatial sensory signals into distinguishable representations that can be retrieved despite noisy changes in background context." (Fyhn et al., 2004).

More recently, a specific evolutionary theory has been put forward regarding the relationship between space and memory in the hippocampal formation (Buzsaki and Moser, 2013). While spatial contextual information has always been considered a critical component of episodic memory, or memory for past events in one's life (Tulving et al., 1972), this theory suggests that spatial and mnemonic representations coexist in medial temporal lobe structures because the mechanisms of memory evolved from mechanisms of navigation (Buzsaki and Moser, 2013). The suggestion is that travel through mental space as one recalls information and previous experiences may recruit the same neural circuitry as navigation through real space. A critical prerequisite for this idea is that spatial representations may reflect locations that the animal is holding in mind, without actually being in that location. This prerequisite has been fulfilled through recordings in the rodent

hippocampus which demonstrate that ensembles of hippocampal place cells fire in sequences that can represent places in front of and behind the animal (Diba and Buzsaki, 2007), providing representations of future possible paths (Johnson and Redish, 2007; Singer et al., 2013) and behavioral trajectories towards a remembered goal (Pfeiffer and Foster, 2013). However, it is as yet unknown whether entorhinal grid cells can similarly represent positions apart from the animal's current location. Our work demonstrated that overt attention in the form of eye movements is sufficient to activate the grid cell network in monkeys (Killian et al., 2012), which suggests that grid cells may be able to support other cognitive functions besides spatial navigation. Accordingly, it is possible that spatial representations in place and grid cells may reflect the location of attention, rather than the location of the animal (Wilming et al., 2014), and importantly, that activation of spatial representations by attention can be directed either externally, to the environment or internally, through mental travel. Further experiments with monkeys, which can be trained to attend to peripheral locations while maintaining fixation on a different location will be informative in distinguishing representations of current body position from representations of mental space.

As mentioned above, research in nonhuman primates presents an opportunity to bridge the gap between the more invasive research of cells and circuits that have been investigated in rodents and complex behaviors that can be measured in humans. In particular, I see three areas in which research in monkeys may be able to take advantage of newly available technical developments to significantly advance our understanding of hippocampal formation function. First, studies with rodents navigating virtual reality environments have identified spatial responses, similar to navigation through real space (Harvey et al., 2009; Dombeck et al., 2010). Monkeys can also be trained to navigate through virtual space via a joystick (Jutras et al., 2014), and the development of novel behavioral tasks, including tasks of spatial memory, coupled with neurophysiological recordings and eye tracking will allow for further identification of spatial representations in the primate as well as what functions these representations might serve.

Second, in order to determine the relationship between ensembles of neurons in the hippocampal formation, either with respect to modules in the entorhinal cortex or sequences of place cells in the hippocampus, one has to be able to record from large numbers of neurons simultaneously. Primate neurophysiologists have to date been limited by the size and quality of chronically implanted devices, as well as the electronics necessary to connect implanted arrays to recording hardware. More pressing, even though there are some methods available for recording from large numbers of neurons on the cortical surface, there are currently no commercially available techniques for recording from large numbers of single neurons in deep brain structures, structures that lie 3–4 cm below the dorsal surface in the monkey brain. However, developing technologies in the fabrication of multi-electrode arrays provide great promise in this regard. Fabrication techniques that involve the application of micro-electro-mechanical systems (MEMS)

lithographic processes to produce dense arrays with arbitrary geometries on flexible polyimide-based films hold particular promise for recording from a large number of channels deep in the nonhuman primate brain (Lewis et al., 2014). Finally, telemetric recording systems have seen a great deal of development over the past decade, and newer systems are capable of transmitting or recording onboard wideband signal from a large number of channels (Maurer et al., 2013). Chronic arrays with hundreds of recording channels along with telemetric recordings provide the potential to examine memory consolidation by recording from the same neural ensembles during sleep and waking periods, implement full body and head movements into virtual reality settings, and even record neural activity from fully freely moving monkeys. Together, these experiments have the potential to dramatically increase our understanding of the primate hippocampal formation and the extent to which similarities of function and representation exist across species.

## A MAP OF COGNITION

Considering the reconciliation of theories regarding the spatial and mnemonic functions of the hippocampal formation, two related ideas have emerged. One idea holds that spatial representations, along with representations of time, provide the organizing structure for experience, and in that vein engender the necessary conditions for memory formation (Eichenbaum, 2014). Experiential content that is encountered close by within the structures of time or space would become associated, consistent with the idea of the hippocampal formation as being predominantly involved in relational learning, i.e., the binding or associating, of representations of multiple distinct elements into a coherent representation in memory (O'Keefe and Nadel, 1978; Cohen and Eichenbaum, 1993). A second, closely related idea is that the mechanisms of memory evolved from mechanisms of navigation in the physical world (Buzsaki and Moser, 2013). According to this idea, path integration-based mechanisms that rely on computations of self-motion relative to environmental landmarks may utilize the same mechanisms as episodic memory formation, the formation of memories that are self-referential. Engaging in mental time travel during the retrieval of episodic memories uses the same neural algorithms as navigating through space via path integration. In contrast, mechanisms that support allocentric or map-based navigation may be similar to mechanisms of semantic memory formation. According to this idea, items that have become associated through the formation of memories may lie in close proximity within some mapping of semantic relatedness. It seems that the field is converging on a reconsideration of the spatial responses in the hippocampus and entorhinal cortex as reflecting one piece of what might be called a "Map of Cognition." Rather than envisioning these neurons as providing a "GPS" for the brain, or a map of only our spatial environment, instead these responses may be seen as contributing to a general organization

of cognition. Apart from time and space, there are likely other dimensions along which cognition may be organized. Time and space present easily quantifiable continuous variables, and neural responses in the hippocampal formation have shown striking representations within these dimensions. To probe the extent to which neurons in the hippocampal formation may show discrete (hippocampus) or periodic/multi-bumped (entorhinal cortex) representations of cognition in general, we need to come up with clever ways to dimensionalize other aspects of the content of our experiences.

## REFERENCES

- Alyan S, McNaughton BL. 1999. Hippocampectomized rats are capable of homing by path integration. *Behav Neurosci* 113:19–31.
- Baddeley AD, Warrington EK. 1970. Amnesia and the distinction between long- and short-term memory. *J Verbal Learn Verbal Behav* 9:176–189.
- Boccarda CN, Sargolini F, Thoresen VH, Solstad T, Witter MP, Moser EI, Moser MB. 2010. Grid cells in pre- and parasubiculum. *Nat Neurosci* 13:987–994.
- Burgess N. 2008. Grid cells and theta as oscillatory interference: Theory and predictions. *Hippocampus* 18:1157–1174.
- Buzsaki G, Moser EI. 2013. Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nat Neurosci* 16:130–138.
- Cohen NJ, Eichenbaum H. 1993. *Memory, Amnesia and the Hippocampal System*. Cambridge, Massachusetts: The MIT Press.
- Diba K, Buzsaki G. 2007. Forward and reverse hippocampal place-cell sequences during ripples. *Nat Neurosci* 10:1241–1242.
- Doeller CF, Barry C, Burgess N. 2010. Evidence for grid cells in a human memory network. *Nature* 463:657–661.
- Dombeck DA, Harvey CD, Tian L, Looger LL, Tank DW. 2010. Functional imaging of hippocampal place cells at cellular resolution during virtual navigation. *Nat Neurosci* 13:1433–1440.
- Drachman DA, Arbib J. 1966. Memory and the hippocampal complex. II. Is memory a multiple process? *Arch Neurol* 15:52–61.
- Eichenbaum H. 2014. Time cells in the hippocampus: A new dimension for mapping memories. *Nat Rev Neurosci* 15:732–744.
- Eichenbaum H, Dusek J, Young B, Bunsey M. 1996. Neural Mechanisms of Declarative Memory. *Cold Spring Harb Symp Quant Biol* 61:197–206.
- Ekstrom AD, Kahana MJ, Caplan JB, Fields TA, Isham EA, Newman EL, Fried I. 2003. Cellular networks underlying human spatial navigation. *Nature* 425:184–188.
- Fyhn M, Hafting T, Witter MP, Moser EI, Moser MB. 2008. Grid cells in mice. *Hippocampus* 18:1230–1238.
- Fyhn M, Molden S, Witter MP, Moser EI, Moser MB. 2004. Spatial representation in the entorhinal cortex. *Science* 305:1258–1264.
- Hafting T, Fyhn M, Molden S, Moser MB, Moser EI. 2005. Microstructure of a spatial map in the entorhinal cortex. *Nature* 436:801–806.
- Harvey CD, Collman F, Dombeck DA, Tank DW. 2009. Intracellular dynamics of hippocampal place cells during virtual navigation. *Nature* 461:941–946.
- Jacobs J, Kahana MJ, Ekstrom AD, Mollison MV, Fried I. 2010. A sense of direction in human entorhinal cortex. *Proc Natl Acad Sci USA* 107:6487–6492.
- Jacobs J, Weidemann CT, Miller JF, Solway A, Burke JF, Wei XX, Suthana N, Sperling MR, Sharan AD, Fried I, Kahana MJ. 2013. Direct recordings of grid-like neuronal activity in human spatial navigation. *Nat Neurosci* 16:1188–1190.
- Jeffery KJ. 2013. Neural odometry: The discrete charm of the entorhinal cortex. *Curr Biol* 23:r204–r206.
- Jenkinson A, Wixted JT, Hopkins RO, Squire LR. 2012. Visual working memory capacity and the medial temporal lobe. *J Neurosci* 32:3584–3589.
- Jensen O, Lisman JE. 2000. Position reconstruction from an ensemble of hippocampal place cells: Contribution of theta phase coding. *J Neurophysiol* 83:2602–2609.
- Johnson A, Redish AD. 2007. Neural ensembles in ca3 transiently encode paths forward of the animal at a decision point. *J Neurosci* 27:12176–12189.
- Jutras MJ, Buffalo EA. 2010. Recognition memory signals in the macaque hippocampus. *Proc Natl Acad Sci USA* 107:401–406.
- Jutras MJ, Fries P, Buffalo EA. 2009. Gamma-band synchronization in the macaque hippocampus and memory formation. *J Neurosci* 29:12521–12531.
- Jutras MJ, Solyst JA, Buffalo EA. 2014. Theta activity in the monkey hippocampus during virtual navigation. Program No. 751.15. 2014 Neuroscience Meeting Planner. Washington, DC: Society for Neuroscience Abstracts, 2014. Online.
- Killian NJ, Jutras MJ, Buffalo EA. 2012. A map of visual space in the primate entorhinal cortex. *Nature*.
- Killian NJ, Potter SM, Buffalo EA. 2013. Encoding of saccade direction in the primate entorhinal cortex. Program No. 769.22. 2013 Neuroscience Meeting Planner. San Diego, CA: Society for Neuroscience Abstracts, 2013. Online.
- Kim S, Sapiurka M, Clark RE, Squire LR. 2013. Contrasting effects on path integration after hippocampal damage in humans and rats. *Proc Natl Acad Sci USA* 110:4732–4737.
- Langston RF, Ainge JA, Couey JJ, Canto CB, Bjerknes TL, Witter MP, Moser EI, Moser MB. 2010. Development of the spatial representation system in the rat. *Science* 328:1576–1580.
- Lever C, Burton S, Jeweewee A, O’Keefe J, Burgess N. 2009. Boundary vector cells in the subiculum of the hippocampal formation. *J Neurosci* 29:9771–9777.
- Lewis CM, Fiedler E, Stieglitz T, Fries P. 2014. Flexible multi-electrode arrays for acute and chronic recordings. Program No. 661.10. 2014 Neuroscience Meeting Planner. Washington, DC: Society for Neuroscience Abstracts, 2014. Online.
- Maaswinkel H, Jarrard LE, Whishaw IQ. 1999. Hippocampectomized rats are impaired in homing by path integration. *Hippocampus* 9:553–561.
- Maurer AP, Thome A, Bohne M, Archibeque MJ, Permenter MR, Vogt JA, Sprawka K, Engle JR, Barnes CA. 2013. Telemetric recordings from temporal lobe of a freely moving primate. Program No. 390.11. 2013 Neuroscience Meeting Planner. San Diego, CA: Society for Neuroscience Abstracts, 2013. Online.
- McNaughton BL, Battaglia FP, Jensen O, Moser EI, Moser MB. 2006. Path integration and the neural basis of the “cognitive map.” *Nat Rev Neurosci* 7:663–678.
- Miller JF, Neufang M, Solway A, Brandt A, Trippel M, Mader I, Hefft S, Merkow M, Polyn SM, Jacobs J, Kahana MJ, Schulze-Bonhage A. 2013. Neural activity in human hippocampal formation reveals the spatial context of retrieved memories. *Science* 342:1111–1114.
- Milner B. 1972. Disorders of learning and memory after temporal lobe lesions in man. *Clin Neurosurg* 19:421–446.
- Mishkin M. 1982. A memory system in the monkey. *Philos Trans R Soc Lond Ser B Biol Sci* 298:85–95.
- Moser EI, Kropff E, Moser MB. 2008. Place cells, grid cells, and the brain’s spatial representation system. *Annu Rev Neurosci* 31:69–89.
- Naya Y, Suzuki WA. 2011. Integrating what and when across the primate medial temporal lobe. *Science* 333:773–776.
- O’Keefe J. 1976. Place units in the hippocampus of the freely moving rat. *Exp Neurol* 51:78–109.

- O'Keefe J, Dostrovsky J. 1971. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res* 34:171–175.
- O'Keefe J, Nadel L. 1978. *The Hippocampus as a Cognitive Map*. Oxford: Oxford University Press.
- Parron C, Save E. 2004. Evidence for entorhinal and parietal cortices involvement in path integration in the rat. *Exp Brain Res* 159:349–359.
- Pfeiffer BE, Foster DJ. 2013. Hippocampal place-cell sequences depict future paths to remembered goals. *Nature* 497:74–79.
- Rolls ET. 1999. Spatial view cells and the representation of place in the primate hippocampus. *Hippocampus* 9:467–480.
- Rolls ET, Miyashita Y, Cahusac PM, Kesner RP, Niki H, Feigenbaum JD, Bach L. 1989. Hippocampal neurons in the monkey with activity related to the place in which a stimulus is shown. *J Neurosci* 9:1835–1845.
- Rolls ET, Robertson RG, Georges-Francois P. 1997. Spatial view cells in the primate hippocampus. *Eur J Neurosci* 9:1789–1794.
- Rutishauser U, Mamelak AN, Schuman EM. 2006. Single-trial learning of novel stimuli by individual neurons of the human hippocampus-amygdala complex. *Neuron* 49:805–813.
- Rutishauser U, Ross IB, Mamelak AN, Schuman EM. 2010. Human memory strength is predicted by theta-frequency phase-locking of single neurons. *Nature* 464:903–907.
- Sargolini F, Fyhn M, Hafting T, McNaughton BL, Witter MP, Moser MB, Moser EI. 2006. Conjunctive representation of position, direction, and velocity in entorhinal cortex. *Science* 312:758–762.
- Sherrill KR, Erdem UM, Ross RS, Brown TI, Hasselmo ME, Stern CE. 2013. Hippocampus and retrosplenial cortex combine path integration signals for successful navigation. *J Neurosci* 33:19304–19313.
- Shrager Y, Kirwan CB, Squire LR. 2008. Neural basis of the cognitive map: Path integration does not require hippocampus or entorhinal cortex. *Proc Natl Acad Sci USA* 105:12034–12038.
- Singer AC, Carr ME, Karlsson MP, Frank LM. 2013. Hippocampal SWR activity predicts correct decisions during the initial learning of an alternation task. *Neuron* 77:1163–1173.
- Solstad T, Boccarda CN, Kropff E, Moser MB, Moser EI. 2008. Representation of geometric borders in the entorhinal cortex. *Science* 322:1865–1868.
- Squire LR, Zola-Morgan S. 1991. The medial temporal lobe memory system. *Science* 253:1380–1386.
- Squire LR, Stark CE, Clark RE. 2004. The medial temporal lobe. *Annu Rev Neurosci* 27:279–306.
- Suzuki WA, Amaral DG. 1994. Perirhinal and parahippocampal cortices of the macaque monkey: Cortical afferents. *J Comp Neurol* 350:497–533.
- Suzuki WA, Eichenbaum H. 2000. The neurophysiology of memory. *Ann N Y Acad Sci* 911:175–191.
- Suzuki WA, Miller EK, Desimone R. 1997. Object and place memory in the macaque entorhinal cortex. *J Neurophysiol* 78:1062–1081.
- Tan HM, Bassett JP, O'Keefe J, Cacucci F, Wills TJ. 2015. The development of the head direction system before eye opening in the rat. *Curr Biol* 25:479–483.
- Taube JS. 1995. Head direction cells recorded in the anterior thalamic nuclei of freely moving rats. *J Neurosci* 15:70–86.
- Taube JS, Muller RU, Ranck JB Jr. 1990. Head-direction cells recorded from the postsubiculum in freely moving rats. I. Description and quantitative analysis. *J Neurosci* 10:420–435.
- Tcheang L, Bulthoff HH, Burgess N. 2011. Visual influence on path integration in darkness indicates a multimodal representation of large-scale space. *Proc Natl Acad Sci USA* 108:1152–1157.
- Tulving E, Donaldson W, Bower GH. United States Office of Naval Research. 1972. *Organization of Memory*. New York: Academic Press. 423 p.
- Welinder PE, Burak Y, Fiete IR. 2008. Grid cells: The position code, neural network models of activity, and the problem of learning. *Hippocampus* 18:1283–1300.
- Wills TJ, Cacucci F, Burgess N, O'Keefe J. 2010. Development of the hippocampal cognitive map in preweanling rats. *Science* 328:1573–1576.
- Wilming N, Konig P, Buffalo EA. 2014. Grid cells reflect the locus of attention, even the absence of movement. Program No. 93.18. 2014 Neuroscience Meeting Planner. Washington, DC: Society for Neuroscience Abstracts, 2014. Online.
- Wirth S, Yanike M, Frank LM, Smith AC, Brown EN, Suzuki WA. 2003. Single neurons in the monkey hippocampus and learning of new associations. *Science* 300:1578–1581.
- Yanike M, Wirth S, Smith AC, Brown EN, Suzuki WA. 2009. Comparison of associative learning-related signals in the macaque perirhinal cortex and hippocampus. *Cereb Cortex* 19:1064–1078.
- Yartsev MM, Witter MP, Ulanovsky N. 2011. Grid cells without theta oscillations in the entorhinal cortex of bats. *Nature* 479:103–107.
- Zola-Morgan S, Squire LR, Mishkin M. 1982. The neuroanatomy of amnesia: Amygdala-hippocampus versus temporal stem. *Science* 218:1337.